arXiv:1004.3964v1 [math.CO] 22 Apr 2010

ON SIMSUN AND DOUBLE SIMSUN PERMUTATIONS AVOIDING A PATTERN OF LENGTH THREE

WAN-CHEN CHUANG, SEN-PENG EU, TUNG-SHAN FU, AND YEH-JONG PAN

Abstract. A permutation $\sigma \in \mathfrak{S}_n$ is simsun if for all k, the subword of σ restricted to $\{1, \ldots, k\}$ does not have three consecutive decreasing elements. The permutation σ is double simsun if both σ and σ^{-1} are simsun. In this paper we present a new bijection between simsun permutations and increasing 1-2 trees, and show a number of interesting consequences of this bijection in the enumeration of pattern-avoiding simsun and double simsun permutations. We also enumerate the double simsun permutations that avoid each pattern of length three.

Contents

1. Introduction 1
1.1. Simsun and double simsun permutations 1
1.2. Pattern-avoiding simsun and double simsun permutations 2
1.3. Increasing 1-2 trees 3
2. A bijection between simsun permutations and increasing 1-2 trees 3
2.1. The bijection ϕ 4
2.2. Finding ϕ^{-1} 5
3. Consequences of the bijection ϕ 6
3.1. Restricted to $\mathcal{RS}_n(213)$ 6
3.2. Restricted to $\mathcal{DRS}_n(213)$ 8
3.3. Restricted to $\mathcal{DRS}_n(132, 213)$ 11
3.4. Restricted to $\mathcal{RS}_n(231)$ 12
3.5. Restricted to $\mathcal{RS}_n(231, 213)$ 13
4. Enumeration of $\mathcal{DRS}_n(\omega)$ for other ω-patterns in $\mathfrak{S}_3$ 13
4.1. Avoiding 132 14
4.2. Avoiding 312/231 15
4.3. Avoiding 321/123 17
5. Remarks 18
References 18

1. Introduction

1.1. Simsun and double simsun permutations. For a permutation $\sigma = \sigma_1 \cdots \sigma_n \in \mathfrak{S}_n$, a *descent* of σ is a pair (σ_i, σ_{i+1}) of adjacent elements with $\sigma_i > \sigma_{i+1}$ $(1 \leq i \leq n-1)$, and a

---

S.-P. Eu is partially supported by National Science Council, Taiwan under grants NSC 98-2115-M-390-002-MY3, T.-S. Fu is partially supported by NSC 97-2115-M-251-001-MY2, Y.-J. Pan is partially supported by NSC 98-2115-M-127-001.

ω	$\lvert\mathcal{RS}_n(\omega)\rvert$	$\lvert\mathcal{DRS}_n(\omega)\rvert$
123	6 (for $n \geq 4$)	2 (for $n \geq 6$)
132	S_n	S_n
213	M_n	S_n
231	M_n	2^{n-1}
312	2^{n-1}	2^{n-1}
321	C_n	C_n

Table 1. The number of simsun and double simsun permutations avoiding a pattern of length 3.

double descent of σ is a triple $(\sigma_i, \sigma_{i+1}, \sigma_{i+2})$ of consecutive elements with $\sigma_i > \sigma_{i+1} > \sigma_{i+2}$ $(1 \leq i \leq n-2)$. The permutation σ is called *simsun* if for all k, the subword of σ restricted to $\{1, \dots, k\}$ (in the order they appear in σ) has no double descents. For example, $\sigma = 24351$ is not simsun since when restricted to $\{1, 2, 3, 4\}$ the subword 2431 of σ contains a double descent 431.

Simsun permutations were named after Rodica Simion and Sheila Sundaram [11]. They are a variant of André permutations of Foata and Schützenberger [4], and are related to the enumeration of the monomials of the *cd*-index of $\mathfrak{S}_n$ (see [5, 6]). Chow and Shiu [1] enumerated simsun permutations by descent, using generating functions. Let $\mathcal{RS}_n$ denote the set of simsun permutations in $\mathfrak{S}_n$. Simion and Sundaram proved that

$$|\mathcal{RS}_n| = E_{n+1}, \tag{1}$$

where E_n is the nth *Euler number*, which also counts the number of permutations $\sigma \in \mathfrak{S}_n$ with the property $\sigma_1 > \sigma_2 < \sigma_3 > \sigma_4 < \cdots$, known as *alternating permutations*.

Inspiring by the notion of double alternating permutations proposed by Stanley [9], we call a permutation $\sigma \in \mathfrak{S}_n$ *double simsun* if both σ and σ^{-1} are simsun. For example, $\sigma = 51324$ is simsun but not double simsun since $\sigma^{-1} = 24351 \notin \mathcal{RS}_5$.

1.2. **Pattern-avoiding simsun and double simsun permutations.** Recently, Deutsch and Elizalde [2] enumerated simsun permutations that avoid a pattern or a set of patterns of length 3. For an integer $t \leq n$, let $\omega = \omega_1 \cdots \omega_t \in \mathfrak{S}_t$. We say that σ contains an *ω-pattern* if there are indices $i_1 < i_2 < \cdots < i_t$ such that $\sigma_{i_j} < \sigma_{i_k}$ if and only if $\omega_j < \omega_k$. Moreover, σ is called *ω-avoiding* if σ contains no ω-patterns. Let $\mathcal{RS}_n(\omega)$ denote the set of ω-avoiding permutations in $\mathcal{RS}_n$. One of Deutsch and Elizalde's results [2] is the complete enumeration of $\mathcal{RS}_n(\omega)$, for any $\omega \in \mathfrak{S}_3$. The counting numbers are listed in the second column in Table 1. Some results involve classical numbers such as Catalan number C_n, Motzkin number M_n, secondary structure number S_n, and Fibonacci number F_n (e.g., $|\mathcal{RS}_n(132, 213)| = F_{n+1}$, see [2]).

In this paper we study the enumeration of pattern-avoiding double simsun permutations. For an $\omega \in \mathfrak{S}_t$, the permutation σ is called *ω-avoiding double simsun* if σ is ω-avoiding simsun and σ^{-1} is simsun. Let $\mathcal{DRS}_n(\omega)$ be the set of ω-avoiding double simsun permutations in $\mathfrak{S}_n$. Note that in this case σ^{-1} is not necessarily ω-avoiding. For example, $231 \in \mathcal{DRS}_3(312)$ since $231 \in \mathcal{RS}_3(312)$ and $(231)^{-1} = 312 \in \mathcal{RS}_n$. One of the main results is the following enumerative hierarchy for restricted simsun permutations

$$\mathcal{DRS}_n(132, 213) \subseteq \mathcal{DRS}_n(213) \subseteq \mathcal{RS}_n(213) \subseteq \mathcal{RS}_n, \tag{2}$$

where $|\mathcal{DRS}_n(132,213)| = F_{n+1}$, $|\mathcal{DRS}_n(213)| = S_n$, and $|\mathcal{RS}_n(213)| = M_n$. In particular, we characterize the permutations in $\mathcal{DRS}_n(213)$ among the permutations in $\mathcal{RS}_n(213)$ by a pattern-condition (Theorem 3.6). Moreover, we give a unified approach to prove these results based on a bijection between simsun permutations and increasing 1-2 trees.

1.3. **Increasing 1-2 trees.**

A rooted tree on the vertex set $[0,n] := \{0,1,\dots,n\}$ is *increasing* if every path from the root is increasing. The vertices with no children are called *leaves*, and the other vertices are called *inner nodes.* Let $\mathcal{T}_n$ denote the set of increasing trees on $[0,n]$ such that every vertex has at most two children. (The order of the subtrees of a vertex is irrelevant.) Members of $\mathcal{T}_n$ are called *increasing 1-2 trees* on $[0,n]$. For example, the five trees in $\mathcal{T}_3$ are shown in Figure 1.



FIGURE 1. The increasing 1-2 trees with four vertices.

Increasing 1-2 trees appeared in connection to the enumeration of alternating permutations (e.g., see [3, 7]). There is a known bijection between increasing 1-2 trees and simsun permutations, due to Maria Monks (mentioned in [10, Solution to Exercise 120]), which is given in terms of flip equivalence classes of increasing binary trees. In this paper we present a new bijection $\phi : \mathcal{T}_n \to \mathcal{RS}_n$ (Theorem 2.1), which has a number of interesting consequences in the enumeration of pattern-avoiding simsun/double simsun permutations. We also enumerate the double simsun permutations that avoid each pattern of length 3.

This paper is organized as follows. The bijection $\phi : \mathcal{T}_n \to \mathcal{RS}_n$ is given in section 2. With the bijection ϕ restricted to (unlabeled) ordered 1-2 trees we enumerate the sets in the hierarchy (2), $\mathcal{RS}_n(231)$, and $\mathcal{RS}_n(231,213)$ in section 3. The enumeration of $\mathcal{DRS}_n(\omega)$, for $\omega \in \{132,312,231,321,123\}$, and $\mathcal{DRS}_n(312,231)$ is given in section 4. In particular, we give simple constructions for the permutations in $\mathcal{DRS}_n(312)$ and $\mathcal{DRS}_n(231)$.

2. A BIJECTION BETWEEN SIMSUN PERMUTATIONS AND INCREASING 1-2 TREES

In this section we prove the following theorem.

Theorem 2.1. *There is a bijection $\phi : \mathcal{T}_n \to \mathcal{RS}_n$ such that a tree $T \in \mathcal{T}_n$ with $k+1$ leaves is carried to a permutation $\phi(T) \in \mathcal{RS}_n$ with k descents.*

Given a $T \in \mathcal{T}_n$, we write T in a *canonical form* such that if a vertex x has two children u, v with $u > v$ then u is the *left child*, v is the *right child.* The vertices u, v are *siblings.* We make the convention that if x has only one child then it is the right child of x. For two vertices $x, y \in T$, we say that y is a *descendant* of x if x is contained in the path from y to the root. Let $\tau(x)$ denote the subtree of T consisting of x and the descendants of x, and let $T - \tau(x)$ denote the subgraph of T when $\tau(x)$ is removed.

2.1. **The bijection ϕ.** Given a set $X \subseteq [n]$ and an increasing 1-2 tree T on $X \cup \{0\}$, we associate T with a word $\phi(T)$ of length $|X|$ with alphabet X and without repeated letters by the following algorithm. By the *inorder* traversal of a tree we mean visiting the left subtree (possibly empty), the root, and then the right subtree, recursively.

Algorithm A.

(A1) If T consists of the root vertex then T is associated with an empty word.

(A2) Otherwise the word $\phi(T)$ is defined inductively by the factorization

$$\phi(T) = \omega \cdot \phi(T'),$$

where the subword ω and the subtree T' are determined as follows.

- If the root of T has only one child x then let $\omega = x$ (consisting of a single letter x) and let $T' = \tau(x)$ (i.e, obtained from T by deleting the root of T), and relabel the vertex x by 0.
- If the root of T has two children u, v with $u > v$ then traverse the left subtree $\tau(u)$ in inorder and write down the word ω of the vertices of $\tau(u)$. Let $T' = T - \tau(u)$ (i.e., obtained by removing $\tau(u)$ from T).

Example 2.2. Let T be the tree shown in Figure 2(a), which is in the canonical form. Since the root of T has two children, the word $\phi(T)$ can be factorized as $\phi(T) = \omega_1 \cdot \phi(T_1)$, where $\omega_1 = 3846$ is the inorder of the vertices of the left subtree $\tau(3)$, and $T_1 = T - \tau(3)$ shown as Figure 2(b). Since the root of T_1 has only one child, $\phi(T_1)$ can be further factorized as $\phi(T_1) = \omega_2 \cdot \phi(T_2)$, where $\omega_2 = 1$ and $T_2 = \tau(1)$ shown as Figure 2(c). Inductively, $\phi(T_2) = 9572$. We then obtain the corresponding word $\phi(T) = 384619572$.



FIGURE 2. The iterative stages of the bijection ϕ for Example 2.2.

Proposition 2.3. *For every $T \in \mathcal{T}_n$, the word $\phi(T)$ is a simsun permutation in $\mathfrak{S}_n$.*

Proof. We shall prove that for all k, the subword ω of $\phi(T)$ restricted to $\{1, \ldots, k\}$ contains no double descents. Note that such a word ω is the word associated with the subtree of T obtained by removing the vertices $n, n-1, \ldots, k+1$ from T. Thus it suffices to prove that $\phi(T)$ contains no double descents for any $T \in \mathcal{T}_n$ and for all n. We proceed by induction on the number of vertices of T. For $n = 1$ and 2, the two cases are trivial. For $n \geq 3$, we distinguish the following two cases.

(i) The root of T has two children. Let u be the left child of the root. Note that the right child of the root must be the vertex 1.) Since ω is the inorder of the vertices in $\tau(u)$, a pair $(\omega_i, \omega_{i+1}) \subseteq \omega \cup \{1\}$ is a descent whenever the vertex ω_i is a leaf in $\tau(u)$. Moreover, within the inorder, there is at least one inner node between any two leaves. It follows that there are no double descents in $\omega \cup \{1\}$. The remaining part of $\phi(T)$ can be checked as in case (ii).

(ii) The root of T has only one child. Let $0 = x_0, x_1, \dots, x_t = x$ be the path that connects the root and the vertex x, where x_j is the only child of x_{j-1} $(1 \le j \le t)$ and x is the first vertex in depth-first-search order of T that has two children. Then $\phi(T)$ can be factorized as $\phi(T) = \mu_1\mu_2$, where $\mu_1 = 1 \cdots t$ is the initial subword and μ_2 is the remaining part. Clearly, μ_1 has no descents. The subword μ_2, which is determined by the subtree $\tau(x)$, can be checked as in case (i).

By induction we prove that $\phi(T)$ contains no double descents. The assertion follows. □

2.2. **Finding ϕ^{-1}.** For a tree $T \in \mathcal{T}_n$ and $i \in [n]$, let $V(i)$ denote the vertex i in T. By the *rightmost path* of T we mean the path from the root to the last vertex in depth-first-search order of T.

Given a $\sigma \in \mathcal{RS}_n$, we shall recover the tree $\phi^{-1}(\sigma)$ by constructing a sequence of trees $T_1, T_2, \dots, T_n = \phi^{-1}(\sigma)$, where T_i is obtained from T_{i-1} by attaching the vertex $V(i)$ to some vertex u of T_{i-1} so that $V(i)$ is a child of u. In fact, T_i corresponds to the subword of σ restricted to $\{1, \dots, i\}$.

Algorithm B.

Initially T_1 is the tree with $V(1)$ attached to the root 0. Suppose we have constructed up to T_{j-1} for some $j \ge 2$. Let $\omega = \omega_1 \cdots \omega_{j-1}$ be the subword of σ restricted to $\{1, \dots, j-1\}$. To construct T_j, we add $V(j)$ to T_{j-1} according to the following cases.

(B1) The element j appears after ω_{j-1} in σ. Then we attach $V(j)$ to the last vertex (in depth-first-search order) of T_{j-1}.

(B2) The element j appears before ω_1 in σ. If the root of T_{j-1} has only one child then we attach $V(j)$ to the root, otherwise we attach $V(j)$ to $V(\omega_1)$.

(B3) The element j is between ω_{i-1} and ω_i in σ, for some $i \le j-1$. There are two cases.
  (a) $\omega_{i-1} > \omega_i$. We attach $V(j)$ to $V(\omega_{i-1})$.
  (b) $\omega_{i-1} < \omega_i$. If $V(\omega_i)$ is in the rightmost path of T_{j-1} then we attach $V(j)$ to $V(\omega_{i-1})$, otherwise we attach $V(j)$ to $V(\omega_i)$.

Example 2.4. Take $\sigma = 53418672 \in \mathcal{RS}_8$. The sequence $T_1, T_2, \dots, T_8$ of trees constructed for $\phi^{-1}(\sigma) = T_8$ is shown in Figure 3. Note that $V(2)$ is attached to $V(1)$ (shown as T_2) since 2 appears after 1 in σ. For T_3, $V(3)$ is attached to the root 0 (shown as T_3) since 3 appears before the subword 12 in σ and the root of T_2 has only one child. For T_4, $V(4)$ is attached to $V(3)$ since 4 appears between 3 and 1. For T_5, $V(5)$ is attached to $V(3)$ since 5 appears before the subword 3412 in σ and the root of T_4 has two children. For T_6, $V(6)$ is attached to $V(1)$ since 6 appears between 1 and 2, and $V(2)$ is in the rightmost path. For T_7, $V(7)$ ia attached to $V(6)$ since 7 appears between 6 and 2. For T_8, $V(8)$ is attached to $V(6)$ since 8 appears between 1 and 6, and $V(6)$ is not in the rightmost path.

Proposition 2.5. *For every $\sigma \in \mathcal{RS}_n$, the tree $\phi^{-1}(\sigma)$ is an increasing 1-2 tree in $\mathcal{T}_n$.*

Proof. Let $T_1, \dots, T_n$ be the sequence of trees constructed for $\phi^{-1}(\sigma) = T_n$. It is clear that these trees are increasing, and that T_j corresponds to the subword μ of σ restricted to $\{1, \dots, j\}$, for all j. We claim that each T_j is an 1-2 trees and $\mu = \phi(T_j)$.

For T_1, it is trivial. Suppose the assertion holds up to T_{j-1}, for some $j \ge 2$. Let $\phi(T_{j-1}) = \omega_1 \cdots \omega_{j-1}$, i.e., the subword of σ restricted to $\{1, \dots, j-1\}$.

For T_j, it suffices to show that the vertex $V(j)$ is attached to a vertex that has at most one child in T_{j-1}. The cases (B1) and (B2) of algorithm B are clear. For case (a) of (B3),



FIGURE 3. The sequence of increasing 1-2 trees for $\sigma = 53418672$.

since (ω_{i-1}, ω_i) is a descent, $V(\omega_{i-1})$ is a leaf in T_{j-1}. For case (b) of (B3), the element j appears between ω_{i-1} and ω_i in σ and $\omega_{i-1} < \omega_i$. By algorithm A and the fact that ω_{i-1} and ω_i are consecutive in $\phi(T_{j-1})$, we observe that if $V(\omega_i)$ is in the rightmost path then $V(\omega_i)$ is the only child of $V(\omega_{i-1})$ in T_{j-1} (otherwise the left child of $V(\omega_{i-1})$ will appear between ω_{i-1} and ω_i in $\phi(T_{j-1})$). Moreover, if $V(\omega_i)$ is not in the rightmost path then $V(\omega_i)$ has at most one child in T_{j-1} (otherwise due to the vertex-inorder of the subtree $\tau(\omega_i)$, the left child of $V(\omega_i)$ will appear between ω_{i-1} and ω_i in $\phi(T_{j-1})$). Hence T_j is 1-2 tree. It is straightforward to show that $\phi(T_j) = \mu$. The proof is completed. □

Note that (ω_i, ω_{i+1}) is a descent $\phi(T)$ if and only if the vertex $V(\omega_i) \in T$ is a leaf other than the last vertex. This completes the proof of Theorem 2.1.

Remarks. The previously known bijection between $\mathcal{T}_n$ and $\mathcal{RS}_n$, given by Maria Monks [10], makes use of flip equivalence classes of increasing binary trees on vertex set $[n+1]$. By her method the permutation that corresponds to a tree $T \in \mathcal{T}_n$ is essentially determined by all the vertices of T but the greatest vertex, while by our method the requested permutation $\phi(T)$ is determined by all of the non-root vertices of T.

3. CONSEQUENCES OF THE BIJECTION ϕ

In this section, with the benefits of the bijection ϕ we enumerate some families of pattern-avoiding simsun and double simsun permutations.

3.1. Restricted to $\mathcal{RS}_n(213)$.

Let $\mathcal{X}_n$ be the set of (unlabeled) ordered 1-2 trees with $n+1$ vertices. (The order of the subtrees of a vertex is significant.) It is known that $|\mathcal{X}_n| = M_n$ is the nth Motzkin number. For convenience, each tree $T \in \mathcal{X}_n$ is uniquely assigned a vertex-labeling by traversing T in *right-to-left preorder* and labeling the vertices from 0 to n. For example, $\mathcal{X}_3$ consists of the first four trees shown in Figure 1, for which the vertices are increasing in right-to-left preorder, while for the fifth one they are not. Clearly, the resulting trees are in the canonical form. The following result is an immediate consequence of the bijection ϕ when the map ϕ restricts to $\mathcal{X}_n$.

Theorem 3.1.

$$|\mathcal{RS}_n(213)| = M_n.$$

Proof. We shall prove that the map ϕ induces a bijection between $\mathcal{X}_n$ and $\mathcal{RS}_n(213)$.

Given a $T \in \mathcal{X}_n$, suppose the permutation $\phi(T)$ contains 213-patterns. Let (x, y, z) be the 213-pattern with the least element x. If there is more than one choice, choose the first one (in lexicographic order). By the vertex-labeling of T, the vertex z is a descendant of y. Moreover, x is the sibling of y if y is in the rightmost path, and x is the left child of y otherwise. In either case the right-to-left preorder y, z, x of these vertices are not increasing, a contradiction. Hence $\phi(T) \in \mathcal{RS}_n(213)$.

On the other hand, given a $\sigma \in \mathcal{RS}_n(213)$, suppose the vertices of $\phi^{-1}(\sigma)$ are not increasing in right-to-left preorder. Let (v, z) be the first pair of consecutive vertices in this order such that $z \geq v + 2$. Let $x = z - 1$ and let y be the parent of z. We observe that x is not a descendant of y, and (x, y, z) forms a 213-pattern in σ, a contradiction. Hence $\phi^{-1}(\sigma) \in \mathcal{X}_n$. The proof is completed. ☐

Let $\mathcal{M}_n$ denote the set of lattice paths, called *Motzkin paths* of length n, from the origin to the point $(n, 0)$ using *up step* $\mathsf{U} = (1, 1)$, *down step* $\mathsf{D} = (1, -1)$, and *level step* $\mathsf{L} = (1, 0)$ that never pass below the x-axis. It is known that $|\mathcal{M}_n| = M_n$. Now we establish a bijection $\chi : \mathcal{X}_n \to \mathcal{M}_n$. For an ordered 1-2 tree T, we associate T with a word $\chi(T)$ of length $|T| - 1$ with alphabet $\{\mathsf{U}, \mathsf{L}, \mathsf{D}\}$ by the following algorithm.

Algorithm C.

(C1) If T consists of the root vertex then T is associated with an empty word.

(C2) Otherwise the word $\chi(T)$ is defined inductively by the following factorization.

- If the root of T has only one child x then let
$$\chi(T) = \mathsf{L} \cdot \chi(T'),$$
where $T' = \tau(x)$ is the subtree of T rooted at x.
- If the root of T has two children u and v, where u (resp. v) is the left (resp. right) child, then let
$$\chi(T) = \mathsf{U} \cdot \chi(T_1) \cdot \mathsf{D} \cdot \chi(T_2),$$
where $T_1 = \tau(v)$ and $T_2 = \tau(u)$ are the right and left subtrees of the root of T, respectively.

Example 3.2. Take the tree T shown in Figure 4(a). Since the root has two children $(u, v) = (5, 1)$, the path $\chi(T)$ can be factorized as $\chi(T) = \mathsf{U}\chi(T_1)\mathsf{D}\chi(T_2)$, where $T_1 = \tau(1)$ and $T_2 = \tau(5)$. Inductively, $\chi(T_1) = \mathsf{UDL}$ and $\chi(T_2) = \mathsf{ULD}$. The corresponding path $\chi(T)$ is shown in Figure 4(b), where the labels of the steps indicate the corresponding vertices in T.



(a) (b)

FIGURE 4. An ordered 1-2 tree and the corresponding Motzkin path.

To find χ^{-1}, given a Motzkin path π, we start with a root vertex 0 and recover the tree $\chi^{-1}(\pi)$, rooted at 0, inductively by a reverse procedure.

Algorithm D.

(D1) If π is empty then we associate π with the root vertex.

(D2) Otherwise the tree $\chi^{-1}(\pi)$ is defined inductively, according to the following cases.

- If π starts with a level step, then we factorize π as $\pi = \mathsf{L} \cdot \pi'$. Attach a vertex, say x, to the root and construct the subtree $\tau(x) = \chi^{-1}(\pi')$, root at x.
- If π starts with an up step, then we factorize π as $\pi = U\pi_1 D\pi_2$, where U is the first step, D is the first down step returning the x-axis, and π_1, π_2 are Motzkin paths of certain length (possibly empty). Attach a vertex u (resp. v) as the left (resp. right) child of the root, and construct the subtrees $\tau(v) = \chi^{-1}(\pi_1)$ rooted at v and $\tau(u) = \chi^{-1}(\pi_2)$ rooted at v.

The bijection $\chi : \mathcal{X}_n \to \mathcal{M}_n$ is established. Hence along with the map in the proof of Theorem 3.1, we have the following result.

Corollary 3.3. *The map $\chi \circ \phi^{-1}$ is a bijection between the two sets $\mathcal{RS}_n(213)$ and $\mathcal{M}_n$.*

3.2. Restricted to $\mathcal{DRS}_n(213)$.

As mentioned by Callan in [8, A004148], the nth secondary structure number S_n coincides with the number of Motzkin paths in $\mathcal{M}_n$ without consecutive up steps (or equivalently, without consecutive down steps). These paths are called *UU-free* (resp. *DD-free*). The map $\chi \circ \phi^{-1}$ established above has the following consequence when restricted to the set $\mathcal{DRS}_n(213)$.

Theorem 3.4.

$$|\mathcal{DRS}_n(213)| = S_n.$$

To prove this theorem, we characterize the permutations in $\mathcal{DRS}_n(213)$ among the permutations in $\mathcal{RS}_n(213)$, and show that the permutations corresponding to the DD-free paths in $\mathcal{M}_n$ have the same characterization.

The following proposition gives a sufficient condition for determining double simsun permutations among simsun permutations.

Proposition 3.5. *For a $\sigma \in \mathcal{RS}_n$, σ^{-1} is simsun if either σ has no 4132-patterns, or every 4132-pattern of σ is contained in a 51342-pattern.*

Proof. Let $\sigma = \sigma_1 \cdots \sigma_n \in \mathcal{RS}_n$ with $\sigma^{-1} \notin \mathcal{RS}_n$. It suffices to prove that σ has a 4132-pattern that is not contained in any 51342-pattern.

Let t be the least integer such that the subword of σ^{-1} restricted to $\{1, \dots, t\}$ contains a double descent $(\sigma_i^{-1}, \sigma_j^{-1}, \sigma_k^{-1})$, where $i < j < k$ and $\sigma_i^{-1} = t$. Let $(\sigma_j^{-1}, \sigma_k^{-1}) = (s, r)$. It follows that $(\sigma_r, \sigma_s, \sigma_t) = (i, j, k)$ is a decreasing triple of σ restricted to $\{1, \dots, k\}$. The relative orders of these elements in σ^{-1} and σ are shown by the diagrams in Figure 5. Since $\sigma \in \mathcal{RS}_n$, $(\sigma_r, \sigma_s, \sigma_t)$ is not a double descent. There are two cases.

(i) There exists an element $\sigma_p = q$ such that $q < j$ and $r < p < s$. Note that if $i < q < j$ then $\sigma_q^{-1} = p < s = \sigma_j^{-1}$, which is against the condition that $(\sigma_i^{-1}, \sigma_j^{-1}, \sigma_k^{-1})$ is a double descent in the subword of σ^{-1} restricted to $\{1, \dots, t\}$. Hence $q < i$, and the quadruple $(\sigma_r, \sigma_p, \sigma_s, \sigma_t)$ is a 4132-pattern in σ (see Figure 5).

(ii) There exists an element $\sigma_p = q$ such that $j < q < k$ and $s < p < t$. Then $\sigma_j^{-1} < \sigma_q^{-1} < t$, which is also against the above condition. This eliminates case (ii).

Moreover, if the 4132-pattern $(\sigma_r, \sigma_p, \sigma_s, \sigma_t)$ is contained in a 51342-pattern then there exists an element $\sigma_g = h$ such that $j < h < k$ and $s < g < t$. This possibility is eliminated as in case (ii). The proof is completed. □



FIGURE 5. The relative order of $(\sigma_r, \sigma_p, \sigma_s, \sigma_t)$ for the permutations σ^{-1} and σ.

Note that a double simsun permutation does not necessarily satisfy the condition in Proposition 3.5. For example, the word 35142 is double simsun with a 4132-pattern 5142 not contained in any 51342-pattern. However, when restricted to 213-avoiding simsun permutations the condition in Proposition 3.5 turns out to be the necessary condition.

Theorem 3.6. *For a $\sigma \in \mathcal{RS}_n(213)$, σ^{-1} is simsun if and only if either σ has no 4132-patterns, or every 4132-pattern of σ is contained in a 51342-pattern.*

Proof. The 'if' part follows from Proposition 3.5. For the 'only if' part, given a $\sigma \in \mathcal{RS}_n(213)$ with $\sigma^{-1} \in \mathcal{RS}_n$, it suffices to prove that if σ has a 4132-pattern, say $(\sigma_m, \sigma_i, \sigma_j, \sigma_k)$, then the quadruple is contained in a 51342-pattern.

Let $(\sigma_m, \sigma_i, \sigma_j, \sigma_k) = (t, z, s, r)$. The relative orders of these elements are shown by the diagrams in Figure 6. Then $(\sigma_r^{-1}, \sigma_s^{-1}, \sigma_t^{-1})$ forms a decreasing triple in the subword of σ^{-1} restricted to $\{1, \dots, k\}$. Since σ^{-1} is simsun, $(\sigma_r^{-1}, \sigma_s^{-1}, \sigma_t^{-1})$ is not a double descent. There are two cases.

(i) There is an element $\sigma_p^{-1} = q$ such that $q < j$ and $r < p < s$. Note that if $q < i$ then $(\sigma_q, \sigma_i, \sigma_j)$ is a 213-pattern in σ, a contradiction. Hence $i < q < j$, and the quintuple $(\sigma_m, \sigma_i, \sigma_q, \sigma_j, \sigma_k)$ forms a 51342-pattern (see Figure 6).

(ii) There is an element $\sigma_p^{-1} = q$ such that $j < q < k$ and $s < p < t$. Then the quintuple $(\sigma_m, \sigma_i, \sigma_j, \sigma_q, \sigma_k)$ forms a 51342-pattern.

In either case the quadruple $(\sigma_m, \sigma_i, \sigma_j, \sigma_k)$ is contained in a 51342-pattern. The proof is completed. □



FIGURE 6. The diagram for a 51342-pattern in a permutation.

In fact, the above characterization for $\mathcal{DRS}_n(213)$ is inspired by the following observation from the map $\chi \circ \phi^{-1} : \mathcal{RS}_n(213) \to \mathcal{M}_n$.

Lemma 3.7. *For a $\sigma \in \mathcal{RS}_n(213)$, the path $\chi \circ \phi^{-1}(\sigma)$ has consecutive down steps if and only if σ has a 4132-pattern that is not contained in any 51342-pattern.*

Proof. Given a $\sigma \in \mathcal{RS}_n(213)$, find the corresponding tree $T = \phi^{-1}(\sigma) \in \mathcal{X}_n$ and path $\pi = \chi(T) \in \mathcal{M}_n$.

Suppose π contains consecutive down steps, say D_2D_1, let U_2 and U_1 be their matching up steps, and let $u_1, v_1, u_2, v_2 \in [n]$ be the vertices in T associated with the steps U_1, D_1, U_2, D_2, accordingly. Then $u_1 > v_1$ are siblings in T, and $u_2 > v_2$ are siblings in the subtree $\tau(v_1)$, as shown in Figure 7. In particular, u_2 must be a leaf since D_2 and D_1 are consecutive in π. By the map ϕ we observe that (u_1, v_1, u_2, v_2) forms a 4132-pattern in permutation σ. Moreover, the quadruple is not contained in any 51342-pattern since u_2 is a leaf in T.

On the other hand, suppose σ has a 4132-pattern that is not contained in any 51342-pattern. Let $(w, x, y, z) \subseteq \sigma$ be the 4132-pattern with the least element w, which is not contained in any 51342-pattern. If there is more than one choice, choose the first one. Then the vertices y, z are descendants of x in T. Moreover, w (resp. y) is the sibling of x (resp. z) if x (resp. z) is in the rightmost path, and w (resp. y) is the left child of x (resp. z) otherwise. In either case w and y are left-child vertices. Let D_1 and D_2 be the down steps in π associated with the siblings of w and y, respectively. Since (w, x, y, z) is not contained in any 51342-pattern, y is a leaf in T, and hence the two down steps D_2 and D_1 are consecutive in π. ☐



FIGURE 7. Two consecutive down steps and the corresponding vertices.

Now we are able to prove Theorem 3.4.

Proof of Theorem 3.4. We shall prove that the map $\chi \circ \phi^{-1}$ in Corollary 3.3 induces a one-to-one correspondence between the permutations in $\mathcal{DRS}_n(213)$ and the DD-free paths in $\mathcal{M}_n$.

Given a $\sigma \in \mathcal{DRS}_n(213)$, let $\pi = \chi \circ \phi^{-1}(\sigma) \in \mathcal{M}_n$. It follows from Theorem 3.6 and Lemma 3.7 that π is DD-free.

On the other hand, given a DD-free path $\pi \in \mathcal{M}_n$, find the corresponding tree $T = \chi^{-1}(\pi) \in \mathcal{X}_n$ and permutation $\sigma = \phi(T) \in \mathcal{RS}_n(213)$. There are two cases.

(i) The height of π is at most 1. Then there are no quadruples $u_1, v_1, u_2, v_2 \in [n]$ such that u_1, v_1 are siblings in T and u_2, v_2 are siblings in the subtree $\tau(v_1)$. Then σ has no 4132-patterns.

(ii) The height of π of is at least 2. By Lemma 3.7, every 4132-pattern in σ is contained in a 51342-pattern.

By Theorem 3.6, σ^{-1} is simsun. Hence $\sigma \in \mathcal{DRS}_n(213)$. The proof is completed. ☐

Next we study the DD-free Motzkin paths that correspond to the permutations in $\mathcal{DRS}_n(132, 213)$.

3.3. **Restricted to $\mathcal{DRS}_n(132, 213)$.** Consider the subset $\mathcal{R}_n \subseteq \mathcal{M}_n$ of Motzkin paths of height 1 with no level steps on the x-axis. For example, $\mathcal{R}_5 = \{\mathsf{ULLLD}, \mathsf{ULDUD}, \mathsf{UDULD}\}$. The nth Fibonacci number F_n also counts the number of *compositions* of $n+1$ with no part equal to 1 (see [8, A000045]). For example, $\{5, 3+2, 2+3\}$ are the requested compositions of 5. It is clear that $|\mathcal{R}_{n+2}| = F_{n+1}$ since the paths in $\mathcal{R}_{n+2}$ are uniquely determined by their block-sizes, which are identical to the compositions of $n+2$.

For a path $\pi \in \mathcal{R}_{n+2}$, we factorize π into blocks as $\pi = \mu_1 \cdots \mu_j$, and then form a new path

$$\pi' = \mu_1 \cdots \mu_{j-1} \mu_j' \tag{3}$$

from π by replacing the last block μ_j by μ_j', where μ_j' is the remaining part of μ_j when the first step and the last step are removed. Note that μ_j' is a segment of level steps (possibly empty) on the x-axis, and $\pi' \in \mathcal{M}_n$. Define $\mathcal{Q}_n = \{\pi' : \pi \in \mathcal{R}_{n+2}\}$. For example, $\mathcal{Q}_3 = \{\mathsf{LLL}, \mathsf{ULD}, \mathsf{UDL}\}$. Clearly, $|\mathcal{Q}_n| = |\mathcal{R}_{n+2}|$.

By the *leftmost path* of an ordered 1-2 tree we mean the maximal path $u_0, u_1, \ldots, u_k$ such that $u_0 = 0$ is the root and u_i is the left child of u_{i-1}, for $1 \le i \le k$. For the paths π' in $\mathcal{Q}_n$, a characterization of the corresponding tree $\chi^{-1}(\pi') \in \mathcal{X}_n$ is that if a vertex x has two children then the vertex x is in the leftmost path.

Theorem 3.8.

$$|\mathcal{DRS}_n(132, 213)| = F_{n+1}.$$

Proof. We shall prove that the map $\chi \circ \phi^{-1}$ induces a bijection between $\mathcal{DRS}_n(132, 213)$ and $\mathcal{Q}_n$.

Given a $\sigma \in \mathcal{DRS}_n(132, 213)$, find the corresponding tree $T = \phi^{-1}(\sigma) \in \mathcal{X}_n$. Suppose there is a vertex x in T with two children $y > z$ such that x is not in the leftmost path. We observe that if x is in the rightmost path then the triple (x, y, z) is a 132-pattern in σ, otherwise the parent of x, say w, together with y and z form a 132-pattern (w, y, z) in σ, which is against the 132-avoiding property of σ. Hence $\chi \circ \phi^{-1}(\sigma) \in \mathcal{Q}_n$.

On the other hand, given a $\pi' \in \mathcal{Q}_n$, find the corresponding tree $T = \chi^{-1}(\pi') \in \mathcal{X}_n$ and permutation $\sigma = \phi(T) \in \mathcal{RS}_n(213)$. Since π' is DD-free, $\sigma \in \mathcal{DRS}_n(213)$. Suppose σ has 132-patterns, let (x, y, z) be the first 132-pattern in σ. Then the vertices y, z are descendants of x in T, and y is either the left child or the sibling of z. In either case there is a vertex with two children in the subtree $\tau(x)$, which is against the characterization of T. Hence $\sigma \in \mathcal{DRS}_n(132, 213)$. □

For the permutations $\sigma \in \mathcal{DRS}_n(132, 213)$, by the above argument, the corresponding trees can be partitioned into paths whose starting points are the ones in the leftmost path. Hence σ can be factorized into maximal increasing subwords (by putting a dot between every descent pair), which can be constructed as follows.

Let $\beta = n(n-1)\cdots 1$ be the word with $\beta_i = n+1-i$, for $1 \le i \le n$. For a composition $C = \{t_1, \ldots, t_j\}$ of n with $t_i \ge 2$ $(2 \le i \le j-1)$ and $t_1, t_j \ge 1$, we factorize β with respect to C as $\beta = \mu_1 \mu_2 \cdots \mu_j$ so that the ith subword μ_i is of length t_i. We associate C with a permutation $\zeta(C) \in \mathcal{DRS}_n(132, 213)$ defined by

$$\zeta(C) = \nu_1 \nu_2 \cdots \nu_j, \tag{4}$$

where ν_i is the word in reverse order of μ_i.

For example, take a composition $C = (1, 3, 2, 3)$ of 9. The factorization of β and the associated permutation $\zeta(C)$ are shown below.

$$C = (1, 3, 2, 3) \quad \longleftrightarrow \quad \beta = 9.876.54.321 \quad \longleftrightarrow \quad \zeta(C) = 967845123.$$

3.4. **Restricted to $\mathcal{RS}_n(231)$.** Based on the bijection $\phi : \mathcal{T}_n \to \mathcal{RS}_n$, we shall establish a connection between $\mathcal{RS}_n(231)$ and $\mathcal{RS}_n(213)$.

Theorem 3.9. *There is a bijection between the two sets $\mathcal{RS}_n(231)$ and $\mathcal{RS}_n(213)$.*

With the map ϕ in Theorem 2.1 we consider the set $\mathcal{F}_n = \{\phi^{-1}(\sigma) | \sigma \in \mathcal{RS}_n(231)\}$ of trees that correspond to 231-avoiding simsun permutations. Note that the vertices of these trees are not necessarily increasing in right-to-left preorder. Moreover, these trees satisfy the condition that if a vertex has two children then its left child is a leaf. With this condition and 231-avoiding property, we establish a bijection $\psi : \mathcal{F}_n \to \mathcal{X}_n$, for which each tree $T \in \mathcal{F}_n$ is uniquely transformed into a tree $\psi(T) \in \mathcal{X}_n$ by the following switching process.

Algorithm E.

(E1) Traverse T in right-to-left preorder, and find the first pair (v, z) of consecutive vertices such that $z \geq v + 2$. Then z is the right child of v (due to 231-avoiding). Moreover, if v has more than one child, let y be the left child of v. One can always find the leaf x with $x = z - 1$ to form a new tree T' by switching the subtrees $\tau(y)$ and $\tau(z)$ from v to x.

(E2) If the vertices of T' are increasing in right-to-left preorder then we are done. Otherwise go to (E1) and proceed to process T'.

Example 3.10. Take a permutation $\sigma = 51324867 \in \mathcal{RS}_8(231)$. The tree $T = \phi^{-1}(\sigma)$ is shown as Figure 8(a). The first consecutive pair (x, v) in right-to-left preorder such that $z \geq v + 2$ is $(v, z) = (2, 4)$. Then T' is obtained by switching $\tau(4)$ to the leaf $x = 3$, shown as Figure 8(b). Repeating the procedure, one obtains the requested tree $\psi(T) \in \mathcal{X}_n$ from T' by switching the subtrees $\tau(8)$ and $\tau(6)$ to the leaf 5, shown as Figure 8(c).



FIGURE 8. Transforming an increasing tree into right-to-left preorder.

To find ψ^{-1}, given a $T \in \mathcal{X}_n$, we shall recover the tree $\psi^{-1}(T) \in \mathcal{F}_n$ by a reverse process.

Algorithm F.

(F1) Traverse T (from left to right) by a depth-first-search, and find the first left-child vertex x such that x is not a leaf. Let w be the sibling of x, and let y, z be the children of x, where y is empty if x has only one child. Find the greatest leaf v in the subtree $\tau(w)$, and form a new tree T' by switching $\tau(y)$ and $\tau(z)$ from x to v.

(F2) If all of the left-child vertices of T' are leaves then we are done. Otherwise go to (F1) and proceed to process T'.

This establishes the bijection $\psi : \mathcal{F}_n \to \mathcal{X}_n$. By Theorem 2.1, we prove that the map $\phi \circ \psi \circ \phi^{-1}$ induces a bijection between the two sets $\mathcal{RS}_n(231)$ and $\mathcal{RS}_n(213)$. This completes the proof of Theorem 3.9.

Moreover, with the map $\chi : \mathcal{X}_n \to \mathcal{M}_n$ the composite $\chi \circ \psi \circ \phi^{-1}$ establishes a connection between the permutations in $\mathcal{RS}_n(231)$ and the paths in $\mathcal{M}_n$.

Corollary 3.11. *The map $\chi \circ \psi \circ \phi^{-1}$ induces a bijection between the two sets $\mathcal{RS}_n(231)$ and $\mathcal{M}_n$.*

Example 3.12. Take again $\sigma = 51324867 \in \mathcal{RS}_8(231)$. As shown in Example 3.10, the tree $\phi^{-1}(\sigma) \in \mathcal{F}_n$ is transformed into a tree $\psi \circ \phi^{-1}(\sigma) \in \mathcal{X}_n$, see Figure 8. Following Example 3.2, we obtain the corresponding Motzkin path $\chi \circ \psi \circ \phi^{-1}(\sigma)$, see Figure 4(b). Note that the number of inversions of σ equals the area under the path $\chi \circ \psi \circ \phi^{-1}(\sigma)$.

Remarks. 1. The bijection $\chi \circ \phi^{-1} : \mathcal{RS}_n(213) \to \mathcal{M}_n$ (in Corollary 3.3) is equivalent to the one given by Deutsch and Elizalde in the third proof of [2, Proposition 4.1], for which the Motzkin paths obtained by their method are in reverse order.

2. The bijection $\chi \circ \psi \circ \phi^{-1} : \mathcal{RS}_n(231) \to \mathcal{M}_n$ (in Corollary 3.11) is equivalent to the one given in the third proof of [2, Proposition 5.1].

3.5. Restricted to $\mathcal{RS}_n(231, 213)$.

We consider the paths that correspond to the fixed points of the map $\psi : \mathcal{F}_n \to \mathcal{X}_n$, i.e., $T = \psi(T)$.

A *weak ascent* in a Motzkin path is a maximal sequence of consecutive up steps and level steps. Let $\mathcal{W}_n \subseteq \mathcal{M}_n$ be the set of Motzkin paths with exactly one weak ascent. For example, $\mathcal{W}_4 = \{\mathsf{LLLL}, \mathsf{LLUD}, \mathsf{LULD}, \mathsf{ULLD}, \mathsf{UUDD}\}$. Note that $|\mathcal{W}_n| = F_{n+1}$, as mentioned by Deutsch in [8, A000045]. For the paths π in $\mathcal{W}_n$, the corresponding trees $\chi^{-1}(\pi) \in \mathcal{X}_n$ can be characterized as ordered 1-2 trees whose left-child vertices are all leaves.

Theorem 3.13.

$$|\mathcal{RS}_n(231, 213)| = F_{n+1}.$$

Proof. We shall prove that map $\chi \circ \phi^{-1}$ induces a bijection between $\mathcal{RS}_n(231, 213)$ and $\mathcal{W}_n$.

Given a $\sigma \in \mathcal{RS}_n(231, 213)$, find the corresponding tree $T = \phi^{-1}(\sigma) \in \mathcal{X}_n$. Suppose T has is a left-child vertex x that is not a leaf, let z be the sibling of x and let y be the right child of x. Then the triple (x, y, z) is a 231-pattern in σ, a contradiction. Hence $\chi \circ \phi^{-1}(\sigma) \in \mathcal{W}_n$.

On the other hand, given a $\pi \in \mathcal{W}_n$, find the corresponding tree $T = \chi^{-1}(\pi) \in \mathcal{X}_n$ and permutation $\sigma = \phi(T) \in \mathcal{RS}_n(213)$. Suppose σ has 231-patterns, let (x, y, z) be the 231-pattern with the least element x. If there is more than one choice, choose the first one. Then the vertex x is either the left child or the sibling of z in T. Moreover, y is the right child of x. Thus x is a left-child vertex with children, which is against the characterization of T. Hence $\sigma \in \mathcal{RS}_n(231, 213)$. ☐

4. ENUMERATION OF $\mathcal{DRS}_n(\omega)$ FOR OTHER ω-PATTERNS IN $\mathfrak{S}_3$

In this section we enumerate pattern-avoiding double simsun permutations for the other patterns of length 3.

4.1. **Avoiding 132.** To enumerate $\mathcal{DRS}_n(132)$, we establish a connection to $\mathcal{DRS}_n(213)$. For this purpose we consider an involution $\Gamma : \mathfrak{S}_n \to \mathfrak{S}_n$ as follows. Let $\sigma = \sigma_1 \cdots \sigma_n \in \mathfrak{S}_n$, and let $\Gamma(\sigma) = \omega_1 \cdots \omega_n$, where ω_i is defined by

$$\omega_i = n + 1 - \sigma^{-1}_{n+1-i}, \tag{5}$$

for $1 \le i \le n$. Note that $\Gamma(\sigma^{-1}) = \Gamma(\sigma)^{-1}$. Let $\mathcal{DRS}_n$ be the set of double simsun permutations in $\mathfrak{S}_n$. For a $\sigma \in \mathcal{DRS}_n$, the following proposition gives a sufficient condition for determining $\Gamma(\sigma)$ being double simsun.

Proposition 4.1. *Given a $\sigma \in \mathcal{DRS}_n$, if $\Gamma(\sigma)$ is not double simsun then $\Gamma(\sigma)$ contains a 42513-pattern, or equivalently, σ contains a 35142-pattern.*

Proof. Let $\sigma = \sigma_1 \cdots \sigma_n \in \mathcal{DRS}_n$, and let $\Gamma(\sigma) = \omega = \omega_1 \cdots \omega_n \notin \mathcal{DRS}_n$. Then either $\omega \notin \mathcal{RS}_n$ or $\omega^{-1} \notin \mathcal{RS}_n$. Suppose $\omega \notin \mathcal{RS}_n$, we claim that ω contains a 42513-pattern.

Let t be the least integer such that the subword of ω restricted to $\{1, \dots, t\}$ contains a double descent $(\omega_i, \omega_j, \omega_k)$, where $i < j < k$ and $\omega_i = t$. Since $\sigma \in \mathcal{DRS}_n$, the following observations hold.

(i) $(j, k) \neq (i + 1, i + 2)$. Otherwise, σ^{-1} contains a double descent

$$\sigma^{-1}_{n+1-k} > \sigma^{-1}_{n+1-j} > \sigma^{-1}_{n+1-i}.$$

(ii) $(\omega_j, \omega_k) \neq (t - 1, t - 2)$. Otherwise, σ contains a double descent

$$\sigma_{n-t+1} > \sigma_{n-t+2} > \sigma_{n-t+3}.$$

It follows from (i) and (ii) that there exists an element ω_m such that $\omega_m > \omega_i$ and $j < m < k$. Now, we consider the following two cases.

Case I. $\omega_j \neq t - 1$. Let $\omega_g = t - 1$. Then ω_g must appear after ω_k, i.e., $g > k$. The reason is that if ω_g appears before ω_j (i.e., $g < j$) then the triple $(\omega_g, \omega_j, \omega_k)$ forms a double descent in the subword of ω restricted to $\{1, \dots, t - 1\}$, which is against the choice of t. Moreover, if ω_g appears between ω_j and ω_k (i.e., $j < g < k$) then this contradicts that $(\omega_i, \omega_j, \omega_k)$ is a double descent in the subword restricted to $\{1, \dots, t\}$. Hence the quintuple $(\omega_i, \omega_j, \omega_m, \omega_k, \omega_g)$ forms a 42513-pattern in ω.

Case II. $\omega_j = t - 1$. Find the greatest element r after ω_j such that $r < \omega_j$, say $r = \omega_g$. Note that $\omega_g \ge \omega_k$. The relative order of $\omega_i, \omega_j, \omega_k$, and ω_g in ω is shown by the diagram on the left of Figure 9. For convenience, let $z' = n + 1 - z$ for every $z \in [n]$. Note that the relative order of the corresponding elements in σ is shown by the diagram on the right of Figure 9, which is obtained from the left one by flipping vertically and then horizontally. We observe that the triple $(\sigma_{n-t+1}, \sigma_{n-t+2}, \sigma_{r'})$ always forms a double descent in the subword of σ restricted to $\{1, \dots, i'\}$. The only concern is that if there exists an element $\sigma_p = q$ such that $j' < q < i'$ and $n - t + 2 < p < r'$ then $(\sigma_{n-t+1}, \sigma_{n-t+2}, \sigma_{r'})$ will no longer be a double descent. However, this will not happen since (ω_i, ω_j) is a descent in the subword of ω restricted to $\{1, \dots, t\}$ and thus there are no such elements $\omega_{q'} = p'$ in ω with $r < p' < t - 1$ and $i < q' < j$. That σ contains a double descent is against the condition that σ is simsun. This eliminates case II.

This proves that if $\omega \notin \mathcal{RS}_n$ then ω contains a 42513-pattern. On the other hand, if $\omega \in \mathcal{RS}_n$ then $\omega^{-1} \notin \mathcal{RS}_n$. By the above argument, ω^{-1} contains a 42513-pattern and so does ω. By the definition of Γ, σ contains a 35142-pattern if and only if ω contains a 42513-pattern. The proof is completed. $\square$



FIGURE 9. The relative order of $(\omega_i, \omega_j, \omega_k, \omega_g)$ for the permutations ω and σ.

Theorem 4.2.

$$|\mathcal{DRS}_n(132)| = S_n.$$

Proof. We shall prove that Γ induces a bijection between $\mathcal{DRS}_n(132)$ and $\mathcal{DRS}_n(213)$.

Given a $\sigma \in \mathcal{DRS}_n(132)$, let $\omega = \Gamma(\sigma)$. Since σ is 132-avoiding, σ contains no 35142-patterns and, by Proposition 4.1, ω is double simsun. Moreover, by the definition of Γ, ω is 213-avoiding. Hence $\omega \in \mathcal{DRS}_n(213)$.

On the other hand, given an $\omega' \in \mathcal{DRS}_n(213)$, let $\sigma' = \Gamma^{-1}(\omega')$. Then $\sigma' = \Gamma(\omega')$. Since ω' is 213-avoiding, ω' contains no 35142-patterns and, by Proposition 4.1, σ' is double simsun. Moreover, by the definition of Γ, σ' is 132-avoiding. Hence $\sigma' \in \mathcal{DRS}_n(132)$.

This proves $|\mathcal{DRS}_n(132)| = |\mathcal{DRS}_n(213)|$. The assertion follows from Theorem 3.4. ☐

Along with the fact $|\mathcal{RS}_n(132)| = S_n$ [2, Theorem 3.1], we prove the relation

$$\mathcal{DRS}_n(132) = \mathcal{RS}_n(132). \tag{6}$$

By the bijection Γ, it is clear that the permutations in $\mathcal{DRS}_n(132)$ and $\mathcal{DRS}_n(213)$ are equidistributed with respect to excedances (i.e., $\sigma_i > i$) and fixed points.

Corollary 4.3. *The number of permutations in $\mathcal{DRS}_n(132)$ with i excedances and j fixed points equals the number of permutations in $\mathcal{DRS}_n(213)$ with i excedances and j fixed points.*

4.2. **Avoiding 312/231.** The following relations show that the sets $\mathcal{RS}_n(312)$, $\mathcal{DRS}_n(312)$, and $\mathcal{DRS}_n(231)$ have the same cardinality.

Lemma 4.4. *The following facts hold.*
- (i) $\mathcal{DRS}_n(312) = \mathcal{RS}_n(312)$.
- (ii) *The map* $\sigma \to \sigma^{-1}$ *is a bijection between* $\mathcal{DRS}_n(312)$ *and* $\mathcal{DRS}_n(231)$.

Proof. (i) It is clear that $\mathcal{DRS}_n(312) \subseteq \mathcal{RS}_n(312)$. On the other hand, we observe that if $\sigma \in \mathcal{RS}_n(312)$ then σ contains no 4132-patterns. It follows from Proposition 3.5 that σ^{-1} is simsun. Hence $\sigma \in \mathcal{DRS}_n(312)$.

(ii) Note that for any $\sigma \in \mathfrak{S}_n$, σ^{-1} is 231-avoiding if and only if σ is 312-avoiding. By definition, it is clear that the map $\sigma \to \sigma^{-1}$ is a bijection between $\mathcal{DRS}_n(231)$ and $\mathcal{DRS}_n(312)$. ☐

Recall that the number 2^{n-1} counts the number of compositions of n. For example, the compositions of 4 consist of $\{4, 3+1, 2+2, 1+3, 2+1+1, 1+2+1, 1+1+2, 1+1+1+1\}$. In the following we present simple constructions for the permutations in $\mathcal{DRS}_n(312)$ and $\mathcal{DRS}_n(231)$, respectively, using compositions of n.

Theorem 4.5.

$$|\mathcal{DRS}_n(231)| = |\mathcal{DRS}_n(312)| = |\mathcal{RS}_n(312)| = 2^{n-1}.$$

Let $\alpha = 1\cdots n$ be the word with $\alpha_i = i$, for $1 \le i \le n$. For a composition $C = (t_1, t_2, \dots, t_j)$ of n, let $s_i = t_1 + \cdots + t_i$, for $1 \le i \le j-1$. For convenience, let $s_0 = 0$ and $s_j = n$. We factorize α with respect to C as $\alpha = \mu_1\mu_2\cdots\mu_j$ by putting a dot between s_i and $s_i + 1$, for $1 \le i \le j-1$. Namely, the ith subword is $\mu_i = (s_{i-1}+1)\cdots s_i$. Then we associate C with the permutation $\rho(C) \in \mathfrak{S}_n$ defined by

$$\rho(C) = \widehat{\mu}_1\widehat{\mu}_2\cdots\widehat{\mu}_j, \tag{7}$$

where $\widehat{\mu}_i = (s_{i-1}+2)\cdots s_i(s_{i-1}+1)$ is the word obtained from μ_i by moving the first element $(s_{i-1}+1)$ to the end of μ_i.

For example, take $C = (3,2,1,3)$, a composition of 9. The factorization of α and the associated permutation $\rho(C)$ are shown below.

$$C = (3,2,1,3) \quad \longleftrightarrow \quad \alpha = 123.45.6.789 \quad \longleftrightarrow \quad \rho(C) = 231546897.$$

Proposition 4.6. *The map ρ is a bijection between the set of compositions of n and the set $\mathcal{RS}_n(312)$.*

Proof. For a composition C of n, it is straightforward to verify that the associated permutation $\rho(C)$ is 312-avoiding simsun.

On the other hand, given a $\sigma \in \mathcal{RS}_n(312)$, we observe that for all k, the subword $\omega_1\cdots\omega_k$ of σ restricted to $\{1,\dots,k\}$ satisfies the condition that either $\omega_{k-1} = k$ or $\omega_k = k$. One can factorize σ into subwords as follows. For $k = 1,\dots,n$, check the subword $\omega_1\cdots\omega_k$ of σ restricted to $\{1,\dots,k\}$, and put a dot in σ at the end of k if $\omega_k = k$. Then the lengths of these subwords form the requested composition $\rho^{-1}(\sigma)$ of n. The assertion follows. □

By Lemma 4.4 and Proposition 4.6, the proof of Theorem 4.5 is completed.

Since $\sigma \in \mathcal{DRS}_n(312) = \mathcal{RS}_n(312)$ if and only if $\sigma^{-1} \in \mathcal{DRS}_n(231)$, we have a similar construction for the permutations in $\mathcal{DRS}_n(231)$.

For a composition $C = (t_1, t_2, \dots, t_j)$ of n and the factorization $\alpha = \mu_1\mu_2\cdots\mu_j$ of α with respect to C, the corresponding permutation $\varrho(C) \in \mathcal{DRS}_n(231)$ is defined by

$$\varrho(C) = \overline{\mu}_1\overline{\mu}_2\cdots\overline{\mu}_j, \tag{8}$$

where $\overline{\mu}_i = s_i(s_{i-1}+1)\cdots(s_i - 1)$ is obtained from μ_i by moving the lase element s_i to the beginning of μ_i. It is clear that $\varrho(C)$ is the inverse of $\rho(C)$. We have the following result.

Proposition 4.7. *The map ϱ is a bijection between the set of compositions of n and the set $\mathcal{DRS}_n(231)$.*

Counting the compositions of n by the number of parts greater than 1 yields the following result.

Corollary 4.8. *The number of permutations in $\mathcal{DRS}_n(312)$, and respectively in $\mathcal{DRS}_n(231)$, with k descents is $\binom{n}{2k}$.*

Proof. Note that the number of compositions C of n with k parts greater than 1 is $\binom{n}{2k}$. By the construction of $\rho(C)$ and $\varrho(C)$, each of these parts contributes exactly one descent to the permutation $\rho(C)$ as well as $\varrho(C)$, and all the other parts contribute fixed points. □

The permutations in $\mathcal{DRS}_n(312)$ and $\mathcal{DRS}_n(231)$ are essentially equidistributed with respect to excedances and fixed points.

Corollary 4.9. *The number of permutations in $\mathcal{DRS}_n(312)$ with i excedances and j fixed points equals the number of permutations in $\mathcal{DRS}_n(231)$ with $n-i-j$ excedances and j fixed points.*

Proof. Given a $\sigma \in \mathcal{DRS}_n(312)$ with i excedances, j fixed points, and k descents, by the map ρ^{-1}, the permutation σ can be factorized as $\sigma = \widehat{\mu}_1\widehat{\mu}_2\dots\widehat{\mu}_{j+k}$. Let $C = \rho^{-1}(\sigma) = (t_1, \dots, t_{j+k})$, where t_d is the length of the dth subword $\widehat{\mu}_d$, for $1 \le d \le j+k$. Note that among these subwords, there are j subwords of length 1 and k subwords of length at least 2. Moreover, each $\widehat{\mu}_d$ contributes exactly $t_d - 1$ excedances to σ. Hence $i + j + k = n$. By the map ϱ, the permutation $\varrho(C) = \overline{\mu}_1\overline{\mu}_2\dots\overline{\mu}_{j+k} \in \mathcal{DRS}_n(231)$ contains exactly k excedances, j fixed points, and k descents since each subword $\overline{\mu}_d$ of length $t_d \ge 2$ contributes one excedance and one descent to $\varrho(C)$. □

By the bijections ρ and ϱ, it is clear that the permutations in $\mathcal{DRS}_n(312, 231)$ are in one-to-one correspondence with the words obtained by factorizing α into subwords of length 1 or 2, and then interchanging the letters in each subword of length 2. This proves the following result.

Corollary 4.10.

$$|\mathcal{DRS}_n(312, 231)| = F_{n+1}.$$

4.3. **Avoiding 321/123.** We make use of the results in [2] to enumerate the sets $\mathcal{DRS}_n(321)$ and $\mathcal{DRS}_n(123)$.

Theorem 4.11. *We have*

(i) $|\mathcal{DRS}_n(321)| = C_n$,

(ii) $|\mathcal{DRS}_4(123)| = 5$, $|\mathcal{DRS}_5(123)| = 3$, *and* $|\mathcal{DRS}_n(123)| = 2$ *for* $n \ge 6$.

Proof. (i) Note that $|\mathcal{RS}_n(321)| = C_n$. By the same argument as in the proof of Lemma 4.4, we have $\mathcal{DRS}_n(321) = \mathcal{RS}_n(321)$.

(ii) Note that $\mathcal{RS}_4(123) = \{3412, 4231, 4132, 3142, 2413, 2143\}$. By the proof of [2, Proposition 2.1], for $n \ge 4$, each permutation $\sigma = \sigma_1 \cdots \sigma_n \in \mathcal{RS}_n(123)$ can produce a unique permutation in $\mathcal{RS}_{n+1}(123)$ by inserting $n+1$ between σ_1 and σ_2 if $\sigma_1 > \sigma_2$ or to the left of σ_1 if $\sigma_1 < \sigma_2$. To determine $\mathcal{DRS}_n(123)$, we shall eliminate those possibilities $\sigma \in \mathcal{RS}_n(123)$ such that σ^{-1} is not simsun.

For $n \ge 4$, if the subword of σ restricted to $\{1, 2, 3, 4\}$ is 4132 then σ^{-1} is not simsun since σ^{-1} contains a double descent $(\sigma_2^{-1}, \sigma_3^{-1}, \sigma_4^{-1})$. In particular, this eliminates 4132 from $\mathcal{RS}_4(123)$. One can check that the others are double simsun. Then $|\mathcal{DRS}_4(123)| = 5$.

Since $\sigma \in \mathcal{DRS}_n(123)$ if and only if $\sigma^{-1} \in \mathcal{DRS}_n(123)$, the last element σ_n of σ is either 1 or 2. In particular, this eliminates 52413 and 25143 from $\mathcal{RS}_5(123)$, and $\mathcal{DRS}_5(123) = \{53412, 45231, 35142\}$. Moreover, for $n \ge 6$ if the subword of σ restricted to $\{1, \dots, 6\}$ is 635142 then σ^{-1} is not simsun since σ^{-1} contains a double descent $(\sigma_4^{-1}, \sigma_5^{-1}, \sigma_6^{-1})$. In particular, this eliminates 635142 from $\mathcal{RS}_6(123)$, and $\mathcal{DRS}_6 = \{563412, 654231\}$. Note that each member of $\mathcal{DRS}_n(123)$ produces a unique permutation in $\mathcal{DRS}_{n+1}(123)$, for $n \ge 6$. The assertion follows. □

5. Remarks

In this paper we enumerate double simsun permutations that avoid a pattern of length 3. However, the total number of double simsun permutations in $\mathfrak{S}_n$ is still unknown, and the initial values 1, 2, 5, 15, 52, 204, 892, 4297 do not match any known integer sequence in Sloane's Encyclopedia [8]. In addition to this enumerative problem, we are also interested in an analogous characterization result for double simsun permutations as Theorem 3.6, i.e., characterize double simsun permutations among simsun permutations by pattern-conditions.

References

[1] C.-O. Chow, W.C. Shiu, Counting simsun permutations by descents, Ann. Combinatorics, to appear.

[2] E. Deutsch, S. Elizalde, Restricted simsun permutations, preprint, `arXiv:0912.1361` (2009).

[3] R. Donaghey, Alternating permutations and binary increasing trees, J. Combin. Theory Ser. A 18 (1975), 141–148.

[4] D. Foata, M.-P. Schützenberger, Nombres d'Euler et permutations alternantes, in *A Survey of Combinatorial Theory*, J.N. Srivistava, et al., eds., North-Holland, Amsterdam, 1973, pp. 173–187; available at `http://www-irma.u-strasbg.fr/~foata/paper/pub18.html`

[5] G. Hetyei, On the *cd*-variation polynomials of André and simsun permutations, Discrete Comp. Geom. 16 (1996), 259–275.

[6] G. Hetyei, E. Reiner, Permutation trees and variation statistics, European J. Combin. 19 (1998) 847–866.

[7] A.G. Kuznetsov, I.M. Pak, A.E. Postnikov, Increasing trees and alternating permutations. (Russian) Uspekhi Mat. Nauk 49 (1994), 79–110; translation in Russian Math. Surveys 49 (1994), 79–114.

[8] N.J.A. Sloane, The On-Line Encyclopedia of Integer Sequences, published electronically at `http://www.research.att.com/~njas/sequences/`.

[9] R.P. Stanley, A survey of alternating permutations, preprint, `arXiv:0912.4240` (2009).

[10] R.P. Stanley, Enumerative Combintorics I, second edition, Chapter 1, preprint (version 2.3.4), available at `http://www-math.mit.edu/~rstan/papers.html`.

[11] S. Sundaram, The homology of partitions with an even number of blocks, J. Algebraic Combin. 4 (1995) 69–92.

Department of Applied Mathematics, National University of Kaohsiung, Kaohsiung 811, Taiwan, ROC
E-mail address: `m0974103@mail.nuk.edu.tw`

Department of Applied Mathematics, National University of Kaohsiung, Kaohsiung 811, Taiwan, ROC
E-mail address: `speu@nuk.edu.tw`

Mathematics Faculty, National Pingtung Institute of Commerce, Pingtung 900, Taiwan, R.O.C
E-mail address: `tsfu@npic.edu.tw`

Department of Computer Science and Information Engineering, Tajen University, Pingtung 907, Taiwan, R.O.C
E-mail address: `yjpan@mail.tajen.edu.tw`